


Cataract Surgery and your Intraocular Lens (IOL) Choices


WHAT ARE CATARACTS?

Cataracts are a clouding of the natural lens in your eye, which makes vision blurry.¹


EYE WITHOUT CATARACT




EYE WITH CATARACT




Cataract Surgery



CATARACT REMOVAL



LENS INSERTION



VISION RESTORED

Recovery after cataract surgery usually occurs within a few days.² See the chart of options below

Your Intraocular Lens (IOL) Choices

Intraocular Lens Choices

Near

Intermediate

Distance

Astigmatism



Monofocal Lens

- Designed to provide clear distance vision
- You will likely still need glasses for reading (and possibly distance vision)³



Astigmatism-Correcting Monofocal Lens

- Corrects both cataracts and astigmatism
- Designed to provide clear distance vision without glasses
- You will likely still need glasses for reading⁴



Next-Generation Trifocal Lens

- Corrects cataracts
- Designed to provide a full range of vision—near, far and in-between⁵
- Not designed to correct for astigmatism



Next-Generation Trifocal Toric Lens

- Corrects cataracts and astigmatism
- Designed to provide a full range of vision—near, far and in-between⁶



Visit www.seethefullpicture.ca for more information

Intraocular Lens Choices

ASTIGMATISM-CORRECTING IOLS

71% of cataract patients have corneal astigmatism, which is a common condition that may produce blurry vision.

What you may see without glasses after you **correct your cataracts and astigmatism together**

What you may see without glasses after you **correct your cataracts**



The simulated images above are for illustration purposes only.

TRIFOCAL IOLS

Trifocal IOLs are designed to provide clear vision without glasses after cataract surgery at near, intermediate and far distances.



The simulated images above are for illustration purposes only.

AcrySof IQ IOL CAUTION Federal law restricts this device to sale by or on the order of a physician. **WARNINGS:** Careful preoperative evaluation and sound clinical judgment should be used by the surgeon to decide the risk/benefit ratio before implanting lens in patient with any of the conditions described in the Directions for Use labeling. **PRECAUTIONS:** Do not sterilize. Do not store over 45°C. Use only sterile irrigating solutions such as BSS+ or BSS+PLUS. Sterile intraocular irrigating solution studies have shown that color vision discrimination is not adversely affected in individuals with the AcrySof Natural IOL and normal color vision. The effect on vision of the AcrySof Natural IOL in subjects with hereditary color vision defects and acquired color vision defects secondary to ocular disease (e.g. glaucoma, diabetic retinopathy, chronic uveitis) and other retinal or optic nerve diseases has not been studied. In previous studies, posterior capsular opacification (PCO) when present was recognized earlier and clinically significant PCO with the AcrySof PreSTOR lenses compared to the monofocal control. **ATTENTION:** Refer to the Physician Labeling for Use for complete indications, warnings and precautions. The long-term effect of filtering by the light and the clinical efficacy of the filtering in the retina have not been conclusively established.

AcrySof IQ Toric IOL CAUTION Federal law restricts this device to sale by or on the order of a physician. **INDICATIONS:** The AcrySof IQ Toric IOLs are intended for primary implantation in the capsular bag of the eye for the visual correction of aphakia and pre-existing corneal astigmatism secondary to a removable cataract. **WARNINGS:** Careful preoperative evaluation and sound clinical judgment should be used by the surgeon to decide the risk/benefit ratio before implanting lens in patient with any of the conditions described in the Directions for Use labeling. **PRECAUTIONS:** As with any surgical procedure, there are risks involved including, but not limited to, infection, inflammation, and secondary surgical intervention. The effect on vision of the AcrySof Natural IOL in subjects with hereditary color vision defects and acquired color vision defects secondary to ocular disease (e.g. glaucoma, diabetic retinopathy, chronic uveitis, and other retinal or optic nerve diseases) has not been studied.

AcrySof IQ PanOptix Presbyopia Correcting Intraocular Lenses Important Product Information **INDICATIONS:** The AcrySof IQ PanOptix Presbyopia Correcting Intraocular Lenses are intended for primary implantation in the capsular bag in the posterior chamber of the eye for the visual correction of aphakia secondary to a removable cataract. **WARNINGS:** Careful preoperative evaluation and sound clinical judgment should be used by the surgeon to decide the risk/benefit ratio before implanting lens in patient with any of the conditions described in the Directions for Use labeling. **PRECAUTIONS:** As with any surgical procedure, there are risks involved including, but not limited to, infection, inflammation, and secondary surgical intervention. The effect on vision of the AcrySof IQ PanOptix IOL in subjects with hereditary color vision defects and acquired color vision defects secondary to ocular disease (e.g. glaucoma, diabetic retinopathy, chronic uveitis, and other retinal or optic nerve diseases) has not been studied.

AcrySof IQ PanOptix Toric Presbyopia Correcting Intraocular Lenses Important Product Information **INDICATIONS:** The AcrySof IQ PanOptix Toric Presbyopia Correcting Intraocular Lenses are intended for primary implantation in the capsular bag in the posterior chamber of the eye for the visual correction of aphakia and pre-existing corneal astigmatism secondary to a removable cataract. **WARNINGS:** Careful preoperative evaluation and sound clinical judgment should be used by the surgeon to decide the risk/benefit ratio before implanting lens in patient with any of the conditions described in the Directions for Use labeling. **PRECAUTIONS:** As with any surgical procedure, there are risks involved including, but not limited to, infection, inflammation, and secondary surgical intervention. The effect on vision of the AcrySof IQ PanOptix IOL in subjects with hereditary color vision defects and acquired color vision defects secondary to ocular disease (e.g. glaucoma, diabetic retinopathy, chronic uveitis, and other retinal or optic nerve diseases) has not been studied.

References: National Eye Institute Staff Facts About Cataract. National Eye Institute. September 2009. https://nei.nih.gov/health/ataract/ataract_facts2. Mayo Clinic Staff. What you can expect. Cataract Surgery. Mayo Clinic. July 9, 2013. <http://www.mayoclinic.org/tests-procedures/ataract-surgery/details/what-you-can-expect/rec-20229550>. Accessed January 27, 2016. AcrySof IQ IOL Directions for Use. Alcon Canada Inc. AcrySof IQ Toric IOL Directions for Use. Alcon Canada Inc. AcrySof IQ PanOptix IOL Directions for Use. Alcon Canada Inc. AcrySof IQ PanOptix Toric IOL Directions for Use. Alcon Canada Inc.

Clear vision without glasses may be possible after cataract surgery